Brain Research, 451 (1988) 408–410
Elsevier

Author Index

Akesson, T.R., Simerly, R.B. and Micevych, P.E., Estrogen-concentrating hypothalamic and limbic neurons project to the medial preoptic nucleus, 381
Almeida, O.F.X., see Mehmanesh, H., 69
Armstrong-James, M. and Fox, K., Evidence for a specific role for cortical NMDA receptors in slow-wave sleep, 189
Arvidsson, J. and Hellstrand, E., A horseradish peroxidase study of the central projections of primary trigeminal neurons innervating the hard palate in the cat, 197

Battenburg, E.L.F., see Lipkin, W.I., 333
Beuche, W., see Hann, P.G., 126
Bischof, H.-J., see Herrmann, K., 43
Blanchard, J.-C., see Böhme, G.A., 309
Bland, B.H., see Konopacki, J., 33
Bloom, F.E., see Lipkin, W.I., 333
Böhme, G.A., Stutzmann, J.-M. and Blanchard, J.-C., Excitatory effects of cholecystokinin in rat hippocampus: pharmacological response compatible with 'central'- or B-type CCK receptors, 309
Braak, E., see Ohm, T.G., 295
Brazhnik, E.S. and Vinogradova, O.S., Modulation of the afferent input to the septal neurons by cholinergic drugs, 1
Brudzynski, S.M., Wu, M. and Mogenson, G.J., Modulation of locomotor activity induced by injections of carbachol into the tegmental pedunculopontine nucleus and adjacent areas in the rat, 119
Buzsaki, G., see Vanderwolf, C.H., 340

Cain, D.P., see Vanderwolf, C.H., 340
Chiu, T.H., see Yu, O., 376
Cho, W.K., see Park, Y., 225
Cooley, R.K., see Vanderwolf, C.H., 340
Cuello, A.C., see Martinez-Murillo, R., 227
Cutlip, A.C., Lenn, N.J. and Wooten, F.G., Behavioral and metabolic alterations in the opiate withdrawal syndrome induced by lesions of fasciculus retroflexus, 54

Danel, N., see Mons, N., 403
D'Angio, M., Serrano, A., Driscoll, P. and Scatton, B., Stressful environmental stimuli increase extracellular DOPAC levels in the prefrontal cortex of hypoemotional (Roman high-avoidance) but not hyperemotional (Roman low-avoidance) rats. An in vivo voltammetric study, 237
Depaulis, A., Pechnick, R.N. and Liebeskind, J.C., Relationship between analgesia and cardiovascular changes induced by electrical stimulation of the mesencephalic periaqueductal gray matter in the rat, 326
Driscoll, P., see D'Angio, M., 237
Duggan, A.W., Hendry, I.A., Morton, C.R., Hutchison, W.D. and Zhao, Z.Q., Cutaneous stimuli releasing immunoreactive substance P in the dorsal horn of the cat, 261
Durant, P.A.C. and Yaksh, T.L., Micturition in the unanesthetized rat: effects of intrathecal capsaicin, *N*-vanillylnonanamide, 6-hydroxydopamine and 5,6-dihydroxytryptamine, 301

El-Yassir, N., see Fleetwood-Walker, S.M., 213
Emson, P.C., see Yamano, M., 179
Enna, S.J., see Scherer, R.W., 361
Etgen, A.M., see Ng Cheong Ton, J.M., 59

Ferkany, J.W., see Scherer, R.W., 361
Fleetwood-Walker, S.M., Hope, P.J., Mitchell, R., El-Yassir, N. and Molony, V., The influence of opioid receptor subtypes on the processing of nociceptive inputs in the spinal dorsal horn of the cat, 213
Fox, K., see Armstrong-James, M., 189
Friede, R.L., see Hann, P.G., 126
Friedemann, M., see Ng Cheong Ton, J.M., 59
Fujita, S., see Miyake, T., 133
Fukuda, M., see Miyake, T., 133
Fukui, H., see Takemua, M., 386
Fukui, M., see Sakata, S., 85
Funakoshi, M., see Ninomiya, Y., 319

Gardner, E.L., see Ng Cheong Ton, J.M., 59
Geffard, M., see Mons, N., 403
Gerhardt, G.A., see Ng Cheong Ton, J.M., 59
Girgis, S., see Yamano, M., 179
Guyenet, P.G., see Sun, M.-K., 345

Hackett, J.T., see Sun, M.-K., 345
Hal, E.D. and Travis, M.A., Inhibition of arachidonic acid-induced vasogenic brain edema by the non-glucocorticoid 21-aminosteroid U74006F, 350
Hallanger, A.E., see Mitani, A., 397
Hann, P.G., Beuche, W., Neumann, U. and Friede, R.L., The rate of Wallerian degeneration in the absence of immunoglobulins. A study in chick and mouse peripheral nerve, 126
Hatori, T., see Miyake, T., 133
Hellstrand, E., see Arvidsson, J., 197
Hendry, I.A., see Duggan, A.W., 261
Henneberry, R.C., see Novelli, A., 205
Herrmann, K. and Bischof, H.-J., The sensitive period for the morphological effects of monocular deprivation in two nuclei of the tectofugal pathway of zebra finches, 43
Herz, A., see Mehmanesh, H., 69
Hillyard, C.J., see Yamano, M., 179
Hirai, K., see Tatsumi, H., 371
Hope, P.J., see Fleetwood-Walker, S.M., 213
Hutchison, W.D., see Duggan, A.W., 261

Ichikawa, M., Plasticity of intra-amygdaloid connections following the denervation of fibers from accessory olfactory bulb to medial amygdaloid nucleus in adult rat: immunohistochemical study of anterogradely transported lectin (*Phaseolus vulgaris* leucoagglutinin), 248
Ishida, M., see Shinozaki, H., 353
Ito, K., see Mitani, A., 397

Jones, B.E. and Webster, H.H., Neurotoxic lesions of the dorsolateral pontomesencephalic tegmentum-cholinergic cell area in the cat. I. Effects upon the cholinergic innervation of the brain, 13

Kameyama, T., see Ukai, M., 357
Kanno, S. and Okada, Y., Laminar distribution of GABA (γ-aminobutyric acid) in the dorsal lateral geniculate nucleus, Area 17 and Area 18 of the visual cortex, and the superior colliculus of the cat, 172
Karbon, E.W., see Scherer, R.W., 361
Kataoka, K., see Mitani, A., 397
Katayama, Y., see Tatsumi, H., 371
Kato, M., see Sakata, S., 85
Kihara, M., see Kubo, T., 366
Kim, K., see Park, Y., 255
Kitamura, R., see Yamamoto, T., 147
Kitamura, T., see Miyake, T., 133
Kiyama, H., see Takemua, M., 386
Kiyomitsu, Y., see Yamamoto, T., 147
Klangkalya, B., Sripairojthikoon, W., Oparil, S. and Wyss, J.M., High NaCl diet increases anterior hypothalamic α_2-adrenoreceptors in SHR, 77
Kolb, B., see Whishaw, I.Q., 97
Konopacki, J., Bland, B.H. and Roth, S.H., Carbachol-induced EEG 'theta' in hippocampal formation slices: evidence for a third generator of theta in CA3c area, 33
Kubo, T. and Kihara, M., *N*-Methyl-D-aspartate receptors mediate tonic vasodepressor control in the caudal ventrolateral medulla of the rat, 366

Lazdunski, M., see Mourre, C., 274
Lenn, N.J., see Cutlip, A.C., 54
Liebeskind, J.C., see Depaulis, A., 326
Lipkin, W.I., Battenburg, E.L.F., Bloom, F.E. and Oldstone, M.B.A., Viral infection of neurons can depress neurotransmitter mRNA levels without histologic injury, 333
Lysko, P.G., see Novelli, A., 205

MacIntyre, I., see Yamano, M., 179
MacRae, P.G., Spirduso, W.W. and Wilcox, R.E., Reaction time and nigrostriatal dopamine function: the effects of age and practice, 139
Martinez-Murillo, R., Semenenko, F. and Cuello, A.C., The origin of tyrosine hydroxylase-immunoreactive fibers in the regions of the nucleus basalis magnocellularis of the rat, 227
Matsuo, R., see Yamamoto, T., 147
McCarley, R.W., see Mitani, A., 397
Mehmanesh, H., Almeida, O.F.X., Nikolarakis, K.E. and Herz, A., Hypothalamic LH-RH release after acute and chronic treatment with morphine studied in a combined in vivo/in vitro model, 69
Micevych, P.E., see Akesson, T.R., 381
Mickle, J.P., see Yates, B.J., 285
Mitani, A., Ito, K., Hallanger, A.E., Wainer, B.H., Kataoka, K. and McCarley, R.W., Cholinergic projections from the laterodorsal and pedunculopontine tegmental nuclei to the pontine gigantocellular tegmental field in the cat, 397
Mitchell, R., see Fleetwood-Walker, S.M., 213
Miyake, T., Hatori, T., Fukuda, M., Kitamura, T. and Fujita, S., Quantitative studies on proliferative changes of reactive astrocytes in mouse cerebral cortex, 133
Mogenson, G.J., see Brudzynski, S.M., 119
Molony, V., see Fleetwood-Walker, S.M., 213
Mons, N., Danel, N. and Geffard, M., Visualization of L-dihydroxyphenylalanine in rat brain by using specific antibodies, 403
Morton, C.R., see Duggan, A.W., 261
Mourre, C., Nehlig, A. and Lazdunski, M., Cerebral glucose utilization after administration of apamin, a toxin active on Ca^{2+}-dependent K^+ channels, 274

Nakayama, S., see Ukai, M., 357
Nehlig, A., see Mourre, C., 274
Neumann, U., see Hann, P.G., 126
Ng Cheong Ton, J.M., Gerhardt, G.A., Friedemann, M., Etgen, A.M., Rose, G.M., Sharpless, N.S. and Gardner, E.L., The effects of Δ^9-tetrahydrocannabinol on potassium-evoked release of dopamine in the rat caudate nucleus: an in vivo electrochemical and in vivo microdialysis, 59
Niijima, K. and Yoshida, M., Activation of mesencephalic dopamine neurons by chemical stimulation of the nucleus tegmenti pedunculopontinus pars compacta, 163
Nikolarakis, K.E., see Mehmanesh, H., 69
Ninomiya, Y. and Funakoshi, M., Amiloride inhibition of responses of rat single chorda tympani fibers to chemical and electrical tongue stimulations, 319
Novelli, A., Reilly, J.A., Lysko, P.G. and Henneberry, R.C., Glutamate becomes neurotoxic via the *N*-methyl-D-aspartate receptor when intracellular energy levels are reduced, 205

Ogawa, N., Sora, Y.H., Saito, M. and Shimazu, T., Hypothalamic regulation of histidyl-proline diketopiperazine binding sites in the rat liver, 115
Ohm, T.G., Braak, E., Probst, A. and Weindl, A., Neuropeptide Y-like immunoreactive neurons in the human olfactory bulb, 295
Okada, Y., see Kanno, S., 172
Oldstone, M.B.A., see Lipkin, W.I., 333
Oparil, S., see Klangkalya, B., 77

Park, S.D., see Park, Y., 255
Park, Y., Park, S.D., Cho, W.K. and Kim, K., Testosterone stimulates LH-RH-like mRNA level in the rat hypothalamus, 255
Pechnick, R.N., see Depaulis, A., 326
Probst, A., see Ohm, T.G., 295

Reilly, J.A., see Novelli, A., 205
Robertson, B., see Vanderwolf, C.H., 340
Rose, G.M., see Ng Cheong Ton, J.M., 59
Rosenberg, H.C., see Yu, O., 376
Roth, S.H., see Konopacki, J., 33

Saito, M., see Ogawa, N., 115
Sakata, S., Shima, F., Kato, M. and Fukui, M., Effects of thalamic parafascicular stimulation on the periaqueductal gray and adjacent reticular formation neurons. A possible contribution to pain control mechanisms, 85
Scatton, B., see D'Angio, M., 237
Scherer, R.W., Karbon, E.W., Ferkany, J.W. and Enna, S.J., Augmentation of neurotransmitter receptor-stimulated cyclic AMP accumulation in rat brain: differentiation between the effects of baclofen and phorbol esters, 361
Semenenko, F., see Martinez-Murillo, R., 227
Serrano, A., see D'Angio, M., 237
Sharpless, N.S., see Ng Cheong-Ton, J.M., 59
Shima, F., see Sakata, S., 85
Shimazu, T., see Ogawa, N., 115
Shinozaki, H. and Ishida, M., Stizolobic acid, a competitive antagonist of the quisqualate-type receptor at the crayfish neuromuscular junction, 353

Simerly, R.B., see Akesson, T.R., 381
Soechting, J.F., Effect of load perturbations on EMG activity and trajectories of pointing movements, 390
Sora, Y.H., see Ogawa, N., 115
Spirduso, W.W., see MacRae, P.G., 139
Sripairojthikoon, W., see Klangkalya, B., 77
Stutzmann, J.-M., see Böhme, G.A., 309
Sun, M.-K., Young, B.S., Hackett, J.T. and Guyenet, P.G., Rostral ventrolateral medullary neurons with intrinsic pacemaker properties are not catecholaminergic, 345

Takemua, M., Kiyama, H., Fukui, H., Tohyama, M. and Wada, M., Autoradiographic visualization in rat brain of receptors for ω-conotoxin GVIA, a newly discovered calcium antagonist, 386
Tatsumi, H., Hirai, K. and Katayama, Y., Measurement of the intracellular calcium concentration in guinea-pig myenteric neurons by using fura-2, 371
Thompson, F.J., see Yates, B.J., 285
Tohyama, M., see Takemua, M., 386
Tohyama, M., see Yamano, M., 179
Travis, M.A., see Hall, E.D., 350

Ukai, M., Nakayama, S. and Kameyama, T., Apomorphine markedly potentiates naltrexone-induced hypodipsia in the rat, 357

Vanderwolf, C.H., Buzsaki, G., Cain, D.P., Cooley, R.K. and Robertson, B., Neocortical and hippocampal electrical activity following decapitation in the rat, 340
Vinogradova, O.S., see Brazhnik, E.S., 1

Wada, M., see Takemua, M., 386
Wainer, B.H., see Mitani, A., 397
Webster, H.H., see Jones, B.E., 13
Weindl, A., see Ohm, T.G., 295
Whishaw, I.Q. and Kolb, B., Sparing of skilled forelimb reaching and corticospinal projections after neonatal motor cortex removal or hemidecortication in the rat: support for the Kennard doctrine, 97
Wilcox, R.E., see MacRae, P.G., 139
Wooten, G.F., see Cutlip, A.C., 54
Wu, M., see Brudzynski, S.M., 119
Wyss, J.M., see Klangkalya, B., 77

Yaksh, T.L., see Durant, P.A.C., 301
Yamamoto, T., Matsuo, R., Kiyomitsu, Y. and Kitamura, R., Taste effects of 'umami' substances in hamsters as studied by electrophysiological and conditioned taste aversion techniques, 147
Yamano, M., Hillyard, C.J., Girgis, S., MacIntyre, I., Emson, P.C. and Tohyama, M., Presence of a substance P-like immunoreactive neurone system from the parabrachial area to the central amygdaloid nucleus of the rat with reference to coexistence with calcitonin gene-related peptide, 179
Yates, B.J., Thompson, F.J. and Mickle, J.P., Responses of spinal cord neurons following stimulation of A femoral-saphenous venous afferent fibers, 285
Yoshida, M., see Niijima, K., 163
Young, B.S., see Sun, M.-K., 345
Yu, O., Chiu, T.H. and Rosenberg, H.C., A comparison of the effects of midazolam and pentobarbital on the dose–response of GABA-gated Cl^- influx in rat brain microsacs, 376

Zhao, Z.Q., see Duggan, A.W., 261

Erratum

Van der Glas, H.W., De Laat, A., Carels, C. and Van Steenberghe, D., Interactive periodontal and acoustic influences on the masseteric post-stimulus electromyographic complex in man, 444 (1988) 284–294.

On p. 287, right-hand column, 20th line, 'If such an acoustic' *should* read 'If a stimulus-related acoustic'.
On p. 291, Table I, heading, 4th column, 'Δ' *should* read 'S'.

Corrigendum

Sharma, T.R., Chan, W.C. and Gintzler, A.R., Effect of chronic naltrexone administration and its withdrawal on the regional activity of neurons that contain norepinephrine, dopamine and serotonin, 442 (1988) 379–386.

On p. 379, abstract, 9th line, the clause ', as well as the mean content of 3-MT in the thalamus,' should be deleted.
On p. 383, left-hand column, 4th and 5th lines from bottom, the words 'as was the mean content of 3-MT in the thalamus' should be deleted.